

A RATIONAL MAP BETWEEN TWO THREEFOLDS

KENICHIRO KIMURA

Let $V_{33} \subset \mathbb{P}^5$ be the threefold defined by the following 2 cubic equations:

$$\begin{aligned} X_0^3 + X_1^3 + X_2^3 + X_3^3 &= 0 \\ X_2^3 + X_3^3 + X_4^3 + X_5^3 &= 0. \end{aligned}$$

Let \tilde{V}_{33} be the blow up of V_{33} along its singular locus. In [12] it is shown that the third betti number of \tilde{V}_{33} is 2 and the L -series $L(H^3(\tilde{V}_{33}), s) = L(f, s)$ with f the newform of weight 4 on $\Gamma_0(9)$. This is also the L -series of a Hecke character of the field $\mathbb{Q}(\sqrt{-3})$.

Let $E \subset \mathbb{P}^2$ be the curve defined by

$$X_0^3 + X_1^3 + X_2^3 = 0.$$

In Remark 4.5 of [4] it is mentioned that there is a piece of the cohomology $H^3(E^3)$ which has the same L -series as that of $H^3(\tilde{V}_{33})$. According to Tate's conjecture there should be a correspondence between V_{33} and E^3 . We give such a correspondence explicitly. Other relatives of these varieties are the fibered squares of the universal curves for $\Gamma(3)$ and $\Gamma_0(9)$. In [8] and in the section 13 of [9] Schoen constructs correspondences between these varieties and E^3 .

We refer the reader to Remark 4.5 of [4] for a list of threefolds which have a two dimensional Galois representation in H^3 whose L -series is associated to an elliptic modular form of weight 4, and known correspondences among those varieties. We include the relevant works in the reference.

Theorem. *There is a dominant rational map from E^3 to V_{33} of degree 3.*

Proof. Consider the affine piece $\{X_3 \neq 0\}$ of V_{33} which is equal to

$$\text{Spec} \mathbb{Q}[X_0, X_1, X_2, X_4, X_5] / (X_0^3 + X_1^3 + X_2^3 + 1, X_2^3 + 1 + X_4^3 + X_5^3).$$

On the other hand, the affine open $(E - \{X_2 = 0\})^3$ of E^3 is given by

$$\text{Spec} \mathbb{Q}[x_1, y_1, x_2, y_2, x_3, y_3] / (x_1^3 + y_1^3 + 1, x_2^3 + y_2^3 + 1, x_3^3 + y_3^3 + 1).$$

Then there is a map from $(E - \{X_2 = 0\})^3$ to V_{33} induced by the ring homomorphism

$$X_0 \mapsto -x_1y_3, X_1 \mapsto -y_1y_3, X_2 \mapsto x_3, X_4 \mapsto -x_2y_3, X_5 \mapsto -y_2y_3.$$

We denote by $\mathbb{Q}(E^3)$ and $\mathbb{Q}(V_{33})$ the function fields of E^3 and of V_{33} respectively. Then one can see that the field $\mathbb{Q}(E^3)$ is generated over $\mathbb{Q}(V_{33})$ by y_3 with the equation $y_3^3 + X_2^3 + 1 = 0$. So this map is dominant of degree 3.

acknowledgment. The author is grateful to the referee for comments on earlier version of this paper which improved the exposition.

REFERENCES

- [1] Ekedahl, T.; van Geemen, B., *An exceptional isomorphism between modular varieties*, Progr. Math. **89** (1991), 51-74.
- [2] van Geemen, B.; Nygaard, N., *On the Geometry and Arithmetic of Some Siegel Modular Threefolds*, J. of Number Theory **53** (1995), 45-87.
- [3] van Geemen, V.; Top, J., *An Isogeny of K3 surfaces*, mathAG/0309272 (2003).
- [4] Hulek, K.; Spandaw, J.; van Geemen, B.; van Straten, D., *The modularity of the Barth-Nieto quintic and its relatives*, Adv. Geom. **1** (2001), 263-289 also available as AG/0010049.
- [5] Livne, R., *Cubic exponential sums and Galois representations*, Contemp. Math. **67** (1987), 247-261.
- [6] Saito, M-H.; Yui, N., *The modularity conjecture for rigid Calabi-Yau threefolds over \mathbb{Q}* , J. Math. Kyoto Univ. **41** (2001), 403-419.
- [7] Schoen, C., *On the geometry of a special determinantal hypersurface associated to the Mumford-Horrocks vector bundle*, J. Reine Angew. Math. **364** (1986), 85-111.
- [8] Schoen, C., *Zero cycles modulo rational equivalence for some varieties over fields of transcendence degree one*, Proc. Sympos. Pure Math. **46, Part 2** (1987), 463-473.
- [9] Schoen, C., *On the computation of the cycle class map for nullhomologous cycles over the algebraic closure of a finite field*, Ann. Sci. Ecole Norm. Sup. (4) **28** (1995), 1-50.
- [10] van Straten, D., *A quintic hypersurface in \mathbb{P}^4 with 130 nodes*, Topology **32** (1993), 857-864.
- [11] Verrill, H., *The L-series of certain rigid Calabi-Yau threefolds*, J. Number Theory **81** (2000), 509-542.
- [12] Werner, J.; van Geemen, B., *New examples of threefolds with $c_1 = 0$* , Math. Z. **203** (1990), 211-225.
- [13] Yui, N., *The arithmetic of certain Calabi-Yau varieties over number fields*, NATO Sci. Ser. C Math. Phys. Sci. **548** (2000), 515-560.

Institute of Mathematics
University of Tsukuba
Tsukuba
305-8571
Japan
kimurak@math.tsukuba.ac.jp